## AP Calculus AB

**Free-Response Questions** 

## **CALCULUS AB**

**SECTION II, Part A** 

Time—30 minutes
2 Questions

A GRAPHING CALCULATOR IS REQUIRED FOR THESE QUESTIONS.

t (seconds)	0	60	90	120	135	150
f(t) (gallons per second)	0	0.1	0.15	0.1	0.05	0

- 1. A customer at a gas station is pumping gasoline into a gas tank. The rate of flow of gasoline is modeled by a differentiable function f, where f(t) is measured in gallons per second and t is measured in seconds since pumping began. Selected values of f(t) are given in the table.
  - (a) Using correct units, interpret the meaning of  $\int_{60}^{135} f(t) dt$  in the context of the problem. Use a right Riemann sum with the three subintervals [60, 90], [90, 120], and [120, 135] to approximate the value of  $\int_{60}^{135} f(t) dt$ .
  - (b) Must there exist a value of c, for 60 < c < 120, such that f'(c) = 0? Justify your answer.
  - (c) The rate of flow of gasoline, in gallons per second, can also be modeled by  $g(t) = \left(\frac{t}{500}\right) \cos\left(\left(\frac{t}{120}\right)^2\right)$  for

 $0 \le t \le 150$ . Using this model, find the average rate of flow of gasoline over the time interval  $0 \le t \le 150$ .

Show the setup for your calculations.

(d) Using the model g defined in part (c), find the value of g'(140). Interpret the meaning of your answer in the context of the problem.

- 2. Stephen swims back and forth along a straight path in a 50-meter-long pool for 90 seconds. Stephen's velocity is modeled by  $v(t) = 2.38e^{-0.02t}\sin\left(\frac{\pi}{56}t\right)$ , where t is measured in seconds and v(t) is measured in meters per second.
  - (a) Find all times t in the interval 0 < t < 90 at which Stephen changes direction. Give a reason for your answer.
  - (b) Find Stephen's acceleration at time t = 60 seconds. Show the setup for your calculations, and indicate units of measure. Is Stephen speeding up or slowing down at time t = 60 seconds? Give a reason for your answer.
  - (c) Find the distance between Stephen's position at time t = 20 seconds and his position at time t = 80 seconds. Show the setup for your calculations.
  - (d) Find the total distance Stephen swims over the time interval  $0 \le t \le 90$  seconds. Show the setup for your calculations.

**END OF PART A** 


## **CALCULUS AB**

**SECTION II, Part B** 

Time—1 hour
4 Questions


NO CALCULATOR IS ALLOWED FOR THESE QUESTIONS.

- 3. A bottle of milk is taken out of a refrigerator and placed in a pan of hot water to be warmed. The increasing function M models the temperature of the milk at time t, where M(t) is measured in degrees Celsius (°C) and t is the number of minutes since the bottle was placed in the pan. M satisfies the differential equation  $\frac{dM}{dt} = \frac{1}{4}(40 M).$  At time t = 0, the temperature of the milk is 5°C. It can be shown that M(t) < 40 for all values of t.
  - (a) A slope field for the differential equation  $\frac{dM}{dt} = \frac{1}{4}(40 M)$  is shown. Sketch the solution curve through the point (0, 5).



- (b) Use the line tangent to the graph of M at t = 0 to approximate M(2), the temperature of the milk at time t = 2 minutes.
- (c) Write an expression for  $\frac{d^2M}{dt^2}$  in terms of M. Use  $\frac{d^2M}{dt^2}$  to determine whether the approximation from part (b) is an underestimate or an overestimate for the actual value of M(2). Give a reason for your answer.
- (d) Use separation of variables to find an expression for M(t), the particular solution to the differential equation  $\frac{dM}{dt} = \frac{1}{4}(40 M)$  with initial condition M(0) = 5.

© 2023 College Board. Visit College Board on the web: collegeboard.org.



- 4. The function f is defined on the closed interval [-2, 8] and satisfies f(2) = 1. The graph of f', the derivative of f, consists of two line segments and a semicircle, as shown in the figure.
  - (a) Does f have a relative minimum, a relative maximum, or neither at x = 6? Give a reason for your answer.
  - (b) On what open intervals, if any, is the graph of f concave down? Give a reason for your answer.
  - (c) Find the value of  $\lim_{x\to 2} \frac{6f(x) 3x}{x^2 5x + 6}$ , or show that it does not exist. Justify your answer.
  - (d) Find the absolute minimum value of f on the closed interval [-2, 8]. Justify your answer.

х	0	2	4	7
f(x)	10	7	4	5
f'(x)	$\frac{3}{2}$	-8	3	6
g(x)	1	2	-3	0
g'(x)	5	4	2	8

- 5. The functions f and g are twice differentiable. The table shown gives values of the functions and their first derivatives at selected values of x.
  - (a) Let h be the function defined by h(x) = f(g(x)). Find h'(7). Show the work that leads to your answer.
  - (b) Let k be a differentiable function such that  $k'(x) = (f(x))^2 \cdot g(x)$ . Is the graph of k concave up or concave down at the point where x = 4? Give a reason for your answer.
  - (c) Let m be the function defined by  $m(x) = 5x^3 + \int_0^x f'(t) dt$ . Find m(2). Show the work that leads to your answer.
  - (d) Is the function m defined in part (c) increasing, decreasing, or neither at x = 2? Justify your answer.

6. Consider the curve given by the equation  $6xy = 2 + y^3$ .

(a) Show that 
$$\frac{dy}{dx} = \frac{2y}{y^2 - 2x}$$
.

- (b) Find the coordinates of a point on the curve at which the line tangent to the curve is horizontal, or explain why no such point exists.
- (c) Find the coordinates of a point on the curve at which the line tangent to the curve is vertical, or explain why no such point exists.
- (d) A particle is moving along the curve. At the instant when the particle is at the point  $\left(\frac{1}{2}, -2\right)$ , its horizontal position is increasing at a rate of  $\frac{dx}{dt} = \frac{2}{3}$  unit per second. What is the value of  $\frac{dy}{dt}$ , the rate of change of the particle's vertical position, at that instant?

**STOP** 

**END OF EXAM**